

# **Towards a Chemical Taxonomy of Comets: A Critical Comparison of Two Infrared Spectroscopic Methods for Quantitative Measurements of Cometary Water.**

Boncho P. Bonev

*Ritter Astrophysical Research Center, Dept. of Physics and Astronomy  
The University of Toledo*

*And*

*Solar System Exploration Division*

*NASA's GSFC*

*Greenbelt, MD 20771*

*U.S.A.*

[bbonev@kuiper.gsfc.nasa.gov](mailto:bbonev@kuiper.gsfc.nasa.gov)

Neil Dello Russo

*Department of Physics*

*Catholic University of America*

*And*

*Solar System Exploration Division*

*NASA's GSFC*

*Greenbelt, MD 20771*

*U.S.A.*

[neil@kuiper.gsfc.nasa.gov](mailto:neil@kuiper.gsfc.nasa.gov)

Michael J. Mumma

*Goddard Center for Astrobiology*

*And*

*Solar System Exploration Division*

*NASA's GSFC*

*Greenbelt, MD 20771*

*U.S.A.*

[mmumma@kuiper.gsfc.nasa.gov](mailto:mmumma@kuiper.gsfc.nasa.gov)

Michael A. DiSanti

*Goddard Center for Astrobiology*

*And*

*Solar System Exploration Division*

*NASA's GSFC*

*Greenbelt, MD 20771*

*U.S.A.*

[disanti@kuiper.gsfc.nasa.gov](mailto:disanti@kuiper.gsfc.nasa.gov)

Erika L. Gibb

*Department of Physics*

*University of Notre Dame*

*Notre Dame, IN 46456*

*U.S.A.*

[egibb@nd.edu](mailto:egibb@nd.edu)

Karen Magee-Sauer

*Department of Chemistry and Physics*

*Rowan University*

Glassboro, NJ 08028  
U.S.A.  
sauer@rowan.edu

An intense cometary bombardment during the early stages of Earth history might have played a major role in delivering the water and organics that subsequently formed the biosphere. Consequently, the chemical composition of the cometary nucleus (and its diversity) is a key issue for astrobiology. While comets are often grouped according to their orbital properties, it is increasingly clear that such groups likely contain comets formed in diverse regions of the proto-planetary disk. In recent years the emergence of accurate abundance measurements has begun to permit a taxonomic classification based on chemistry.

H<sub>2</sub>O is the principal parent volatile in comets and its sublimation controls the release of other volatiles within 3-4 AU from the Sun. Accurate quantitative production rates of water are critical because they provide the "baseline" for comparison with organic species; their simultaneous measurement eliminates many sources of systematic error. Two methods for deriving cometary water production rates from ground-based high-resolution infrared spectra have now been developed. The H<sub>2</sub>O population can be directly sampled through "hot-band" fluorescent emission [Dello Russo et al. 2000, Icarus]. More recently, vibrational prompt emission from OH has also been used as a proxy for water production [Bonev et al. 2004, ApJ]. In this presentation we critically review the two methods using specific examples from recently analyzed comet data.

This work was supported by grants to MJM (RTOP 344-32-51-03) under NASA's Planetary Astronomy Program, and to NDR (NNG05GA64G) under NASA's Planetary Atmospheres Program.